

03/2020

Press Release

**Summary:**

- The rhizosphere is home to a rich microbial diversity. The metabolites secreted by the roots (products of root exudation) are known to shape the composition of the root microbiota. However, until recently it was not known if or how the microbiota in turn impact root exudation.
- Scientists have discovered that microbial communities can effect specific systemic changes in tomato root exudation via root-to-root signalling.
- The underlying process was termed as “systematically induced root exudation of metabolites” (SIREM).
- The findings were published in PNAS Plant Biology.

## Bacterial influencers – Rhizosphere microbiome mediates root metabolite exudation

Gatersleben, 07.02.2020 **Roots are plant organs, that typically absorb water and minerals from soil. It is lesser known that roots also secrete metabolites, so-called root exudates, which impact the properties of soil directly around the root. This thin layer of soil is called the rhizosphere and is home to a rich microbial diversity, the root microbiota. By producing certain exudates, plants communicate with and govern the microbial life within their rhizosphere for their own benefit. Now, researchers have discovered that this is not a one-way process. Whilst investigating tomato plants, they found that the microbiota can also systemically shape and control root exudation.**

When thinking of ecological hotspots, roots and the earth surrounding them might not immediately spring to mind. However, precisely this region, the rhizosphere, is considered as one of the most complex ecosystems found on earth. It harbours a diverse microbiotic community, including numerous bacteria, fungi and archaea, thriving in an environment rich in biochemical compounds, that are exuded by plant roots at the core of the rhizosphere.


Plants govern the rhizosphere microbiota and shape the soil physical and chemical properties through their root exudates. At the same time, it is well known that roots sense changes in the rhizosphere and trigger systemic responses to defend against pathogens or to adjust to changes in nutrient availability. Nonetheless, there are still many open questions regarding the dynamics and impact of the microbiota on the root itself, and it was not clear how, or whether at all, the rhizosphere microbiota affects the root exudation. An international research team led by Dr. Elisa Korenblum, a scientist from the Weizmann Institute of Science in Israel in collaboration with Dr. Jędrzej Szymanski from the Leibniz Institute IPK in Gatersleben, recently took on this question whilst investigating roots of tomato plants.

Dr Korenblum and her team conducted and analysed a row of split-root experiments, where half of roots of each plant was exposed to a microbiome-rich soil, and the other half was grown in sterile and biochemically ambient conditions. This enabled them to investigate the effect of different microbial communities on the local root system, as well as the systemic changes in the distant roots in anticipation of the presence of new microorganisms. Dr. Szymanski, head of the Network Analysis and Modelling group, traced the complex network of biochemical and gene expression signals controlling this microbiome-root communication and their propagation from the place of origin to distant roots. They thereby discovered that the tomato rhizosphere microbiome can directly affect the chemical composition of roots and root exudates via a systemic root-to-root signalling mechanism. For example, bacteria of the genus *Bacillus* use this process, which the scientists termed

Systemically Induced Root Exudation of Metabolites (SIREM), to trigger the secretion of acylsugars in the whole root system.

The discovery of SIREM is a first step towards untangling the regulatory network spanning the complex plant root–microbial relationship. It is likely that the SIREM process is a key feature of root-microbiota interactions within the rhizosphere, and that the microbiome-reprogrammed systemic root exudation promotes soil conditioning. The precise extent of the regulatory role and incidence of SIREM is yet to be determined.

**Figure (for free use):**



**Illustration:** Bacterial communities trigger systemic signals that lead to regulatory and metabolic changes in distant roots, as well as in green parts of the plant. One function of such signaling is root system-wide management of the rhizosphere microbiota via exudation of specific molecules to the soil. (J. Szymanski / IPK)

**Original publication:** Elisa Korenblum *et. al* (2020) “*Rhizosphere microbiome mediates systemic root metabolite exudation by root-to-root signaling*” PNAS first published February 3, 2020 <https://doi.org/10.1073/pnas.1912130117>

#### Scientific Contact

Dr. Jędrzej Szymanski  
Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben  
Tel.: +49 39482 5753  
E-mail: [szymanski@ipk-gatersleben.de](mailto:szymanski@ipk-gatersleben.de)

Dr. Elisa Korenblum  
Prof. Asaph Aharoni Lab, Weizmann Institute of Science, Rehovot, Israel  
Tel.: +972-8-934-2275  
E-Mail: [elisa.korenblum@weizmann.ac.il](mailto:elisa.korenblum@weizmann.ac.il)

#### Media Contact

Managing Office | Public Relations  
Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben  
Tel. +49 39482 5424  
E-Mail: [geschaefsstelle@ipk-gatersleben.de](mailto:geschaefsstelle@ipk-gatersleben.de)

Weizmann Institute of Science  
234 Herzl Street, Rehovot 7610001, Israel  
Tel: +972 8 934 3852/3856  
E-Mail: [news@weizmann.ac.il](mailto:news@weizmann.ac.il)