

Joint press release of the IPK Leibniz Institute and the Leibniz University  
Hannover

## Researchers discover key gene for toxic alkaloid in barley

Gatersleben, 28.03.2024 **Barley is one of the most important cereal crops on a global scale. Many barley cultivars produce a toxic alkaloid called gramine that affects the suitability of barley as fodder, but also helps to protect barley from pathogens. So far, the potential of manipulating gramine levels has not been harnessed for plant breeding, because the genetic basis of gramine production has been unresolved. Research groups from IPK Leibniz Institute and the Leibniz University Hannover now disclose the complete biosynthetic pathway of gramine and demonstrate how gramine biosynthesis can be introduced into model organisms or removed from barley. Today, the results were published in the journal "Science".**

All plants mediate their environmental interactions via chemical signals. An example is the alkaloid gramine produced by barley, one of the world's most widely grown cereals. Gramine provides protection against herbivorous insects and grazing animals and inhibits the growth of other plants. Despite long-standing interest, the key gene for the formation of gramine remained elusive.

The researchers discovered a cluster of two genes in barley for gramine biosynthesis. The first gene (HvNMT) had already been discovered 18 years ago. In their study the researchers from IPK and the Leibniz University Hannover now identified a second gene (AMI synthase, HvAMIS), and found out that both genes are located in proximity of each other on the same chromosome. With this discovery, the pathway of gramine biosynthesis is now fully elucidated.

"We discovered that AMIS is an oxidase enzyme that carries out an unusual cryptic oxidative rearrangement of tryptophan, allowing us to revise the previous biosynthetic proposal from the 1960s", says Dr. John D'Auria, head of IPK's research group "Metabolic Diversity". Prof. Dr. Jakob Franke, head of the group "Biochemistry of Plant Specialised Metabolites" at Leibniz University Hannover, adds: "We were very surprised by the so far unknown enzyme mechanism by which gramine is formed. At the same time, we now have the possibility to produce biologically active alkaloids with sustainable biotechnological methods."

The research team could produce gramine in yeast and model plants (*Nicotiana benthamiana*, *Arabidopsis*). "In contrast to many other protective metabolites from plants, production of gramine requires only two genes. Therefore, using our findings for practical applications is relatively straightforward", emphasises Ling Chuang from Leibniz University Hannover, one of the first authors. "Furthermore, genetic engineering of barley allowed us to produce gramine in a non-gramine producing barley variety, and eliminate gramine production in a gramine producing barley variety by genome editing", explains the other first author Sara Leite Dias, International Max Planck Research School funded researcher at the IPK.

## Press Release

### Scientific Contact

Dr. John D'Auria  
Phone: +49 39482 5176  
[dauria@ipk-gatersleben.de](mailto:dauria@ipk-gatersleben.de)

Prof. Dr. Jakob Franke  
Phone: +49 511 762 2628  
[jakob.franke@botanik.uni-hannover.de](mailto:jakob.franke@botanik.uni-hannover.de)

### Media Contact

Christian Schafmeister  
Phone: +49 39482 5461  
[schafmeister@ipk-gatersleben.de](mailto:schafmeister@ipk-gatersleben.de)


Mechtild von Münchhausen  
Phone: +49 511 762 3342  
[kommunikation@zuv.uni-hannover.de](mailto:kommunikation@zuv.uni-hannover.de)

“The results set the basis to produce gramine in organisms without the native ability to synthesize it for purposes such as a natural plant protection agent, or to eliminate gramine from barley and other grasses to reduce toxicity towards ruminants”, says Dr. John D’Auria. “Our findings set the ground for improving barley to increase its resistance to pests, reduce its toxicity to ruminants and contribute to sustainable weed management.”

**Original publication:**

Leite Dias *et al.* (2024): Biosynthesis of the allelopathic alkaloid gramine in barley by a cryptic oxidative rearrangement. Science. DOI: [10.1126/science.adk6112](https://doi.org/10.1126/science.adk6112)

**Figure:**



Gramine is a defensive alkaloid from barley (*Hordeum vulgare*) and other Poaceae members. Gramine conveys desirable as well as undesirable traits for breeding. In their study the researchers observed that the formation of gramine and AMI correlates with a distantly clustered cytochrome P450 monooxygenase gene in barley.

The figure shows the loci of HvAMIS and HvNMT on chromosome 1H of the barley variety HOR10350 and the pathway of biosynthesis of gramine which is derived from tryptophan and produced by the now characterized cytochrome P450 monooxygenase that was named AMI Synthase (HvAMIS).